


FP



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) EP 1 043 395 A1

(12) **EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 158(3) EPC

- |                                                                                                                                                    |                                                                                                                                                                                                                                                                                                                              |
|----------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>(43) Date of publication:<br/>11.10.2000 Bulletin 2000/41</p> <p>(21) Application number: 98961610.7</p> <p>(22) Date of filing: 25.12.1998</p> | <p>(51) Int. Cl.<sup>7</sup>: <b>C12N 15/12</b>, C12N 15/63,<br/>C12Q 1/02, C12P 21/08,<br/>C07K 14/705, C07K 16/28,<br/>C12N 1/21, C12N 1/19,<br/>C12N 5/12</p> <p>(86) International application number:<br/>PCT/JP98/05967</p> <p>(87) International publication number:<br/>WO 99/33978 (08.07.1999 Gazette 1999/27)</p> |
|----------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

- (84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**
- (30) Priority: 26.12.1997 JP 36118797
- (71) Applicant:  
**BANYU PHARMACEUTICAL CO., LTD.**  
Chuo-ku, Tokyo 103-8416 (JP)
- (72) Inventors:  
• **ITADANI, Hiraku,**  
Banyu Pharmaceutical Co., Ltd  
Tsukuba-shi, Ibaraki 300-2611 (JP)

- **TAKIMURA, Tetsuo,**  
Banyu Pharmaceutical Co., Ltd  
Tsukuba-shi, Ibaraki 300-2611 (JP)
- **NAKAMURA, Takao,**  
Banyu Pharmaceutical Co., Ltd  
Tsukuba-shi, Ibaraki 300-2611 (JP)
- **OHTA, Masataka,**  
Banyu Pharmaceutical Co., Ltd  
Tsukuba-shi, Ibaraki 300-2611 (JP)
- (74) Representative:  
**VOSSIUS & PARTNER**  
Siebertstrasse 4  
81675 München (DE)

(54) **NOVEL GUANOSINE TRIPHOSPHATE (GTP)-BINDING PROTEIN CONJUGATE TYPE RECEPTOR PROTEINS**

(57) The present invention provides a full-length cDNA encoding a rat G protein-coupled receptor protein, which was isolated by screening cDNA libraries originating from rat thalamus and hypothalamus. It also provides human cDNA corresponding to the rat cDNA. Use of these G protein-coupled receptor proteins makes it possible to screen ligands and also drug-candidate compounds capable of regulating signal transduction from receptors.

EP 1 043 395 A1

**Description**Technical Field

- 5 [0001] The present invention relates to a novel guanosine triphosphate binding protein-coupled receptor protein, a DNA encoding said protein, and a method for screening drug-candidate compounds using them.

Background Art

- 10 [0002] Many hormones and neurotransmitters regulate physiological functions through specific receptor proteins located on the cell membrane. Many of these receptor-proteins transduce signals into the cell by activating a guanosine triphosphate binding protein (occasionally referred to as "G protein" below) that is coupled to them. These receptor proteins are thereby named as G protein-coupled receptors. Since they have a common structure composed of seven transmembrane regions, they are also generally called "seven-transmembrane receptor proteins."

- 15 [0003] G protein-coupled receptors, which are expressed on the surface of cells *in vivo* and functioning cells of tissues, play an extremely important role as a target of molecules such as hormones, neurotransmitters, and biologically active compounds, which regulate the functions of these cells and tissues. Therefore, G protein-coupled receptor proteins have received great attention as targets in drug-development.

- [0004] G protein-coupled receptors reported so far include, muscarinic acetylcholine receptors M1, M2, M3, and M4 (Peralta E.G. et al., EMBO J. 6:3923-3929 (1987)), muscarinic acetylcholine receptor M5 (Bonnér T. I. et al., Neuron 1:403-410 (1988)), adenosine receptor A1 (Libert F. et al., Science 244:569-572 (1989)),  $\alpha 1$ A adrenoreceptor (Bruno J.F. et al., Biochem. Biophys. Res. Commun. 179:1485-1490 (1991)),  $\beta 1$  adrenoreceptor (Friele T. et al., Proc. Natl. Acad. Sci. USA 84:7920-7924 (1987)), angiotensin receptor AT<sub>1</sub> (Takayanagi R. et al., Biochem. Biophys. Res. Commun. 183:910-916 (1992)), endothelin receptor ET<sub>A</sub> (Adachi M. et al., Biochem. Biophys. Res. Commun. 180:1265-1272 (1991)), gonadotropin releasing factor receptor (Kaker S.S. et al., Biochem. Biophys. Res. Commun. 189:289-295 (1992)), histamine receptor H<sub>2</sub> (Ruat M. et al., Proc. Natl. Acad. Sci. USA 87:1658-1672 (1992)), neuropeptide Y receptor Y1 (Larhammar D. et al., J. Biol. Chem. 267:10935-10938 (1992)), interleukin-8 receptor IL8R<sub>A</sub> (Holmes W.E. et al., Science 256:1278-1280 (1991)), dopamine receptor D<sub>1</sub> (Mahan L.C. et al., Proc. Natl. Acad. Sci. USA 87:2196-2200 (1990)), metabolic glutamate receptor mGluR1 (Masu M. et al., Nature 349:760-765 (1991)), and somatostatin receptor SS<sub>1</sub> (Yamada Y. et al., Proc. Natl. Acad. Sci. USA 89:251-255) (for reference, Watson S. and Arkinstall S., The G protein Linked Receptor FactsBook, Academic Press (1994)). Examples of developed medicines aimed at G protein-coupled receptors are: terazosine hydrochloride (antihypertensive agent,  $\alpha 1$  adrenoreceptor antagonist), atenolol (antiarrhythmia,  $\beta 1$  adrenoreceptor antagonist), dicyclomine hydrochloride (antispasmodic agent, acetylcholine receptor antagonist), ranitidine hydrochloride (drug for peptic ulcers, histamine receptor H<sub>2</sub> antagonist), trazodone hydrochloride (antidepressant, serotonin receptor 5-HT<sub>1B</sub> antagonist), and buprenorphine hydrochloride (analgesic agent, opioid receptor  $\kappa$  agonist) (for reference, Stadel J.M. et al., Trends Pharm. Sci. 18:430-437 (1997); Medicine Handbook 5<sup>th</sup> edition, Yakugyo-Jiho).

- [0005] The hypothalamus, a part of the brain which governs a number of programs that trigger a particular response, contributes to the homeostasis of the internal environment by means of a variety of outputs, as the center of the autonomic nervous system. For instance, it releases hormones such as thyrotropic hormone releasing hormone, gonadotropic hormone releasing hormone, and growth hormone releasing hormone, and thereby regulates the entire endocrine system through the actions of these hormones on the specific receptors expressed in target cells. These outputs in the hypothalamus are thought to be mediated by receptors expressed in the hypothalamus and compounds reacting with them. Therefore, elucidation of the relationship between the compounds regulating the hypothalamus outputs and their specific receptors expressed in the hypothalamus is extremely important in developing novel medicines for the treatment of diseases arising from endocrine disorders.

Disclosure of the Invention

- 50 [0006] The present invention provides a novel G protein-coupled receptor protein expressed in the brain (in particular, thalamus and hypothalamus, etc.). It also provides a method for screening ligands and drug-candidate compounds using said receptor protein.

- [0007] The inventors first selected a region highly conserved in known G protein-coupled receptor proteins, then designed primers corresponding to the region, and performed reverse transcriptase-polymerase chain reaction (RT-PCR) using mRNA obtained from rat thalamus and hypothalamus. Next, amplified clones were randomly selected, and their partial nucleotide sequences were determined. To remove known clones from the nucleotide sequence determined-clones, colony-hybridization was performed using as a probe, cDNA clones judged to be encoding a known G protein-coupled receptor protein by homology search. Negative clones that failed to hybridize with any probe were

selected. Using probes prepared based on the nucleotide sequence of the negative clones, the inventors screened cDNA libraries from rat thalamus and hypothalamus, and succeeded in isolating a full-length cDNA encoding a rat G protein-coupled receptor. They also succeeded in isolating a full-length human cDNA corresponding to the rat cDNA. Furthermore, northern blot analysis of the tissue specificity of the gene expression showed that these genes are specifically expressed in the brain.

**[0008]** These G protein-coupled receptors would be extremely useful in screening for ligands and compounds that regulate the signal transduction from receptors, which are anticipated to be utilized as novel medicines.

**[0009]** Thus, the present invention relates to novel G protein-coupled receptor proteins that are expressed in the brain, DNAs encoding them, and a method for screening ligands and compounds that are drug-candidate compounds using the proteins.

**[0010]** Specifically, the invention relates to:

- (1) a guanosine triphosphate binding protein-coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 1, or said amino acid sequence in which one or more amino acids are replaced, deleted, or added;
- (2) a guanosine triphosphate binding protein-coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 20, or said amino acid sequence in which one or more amino acids are replaced, deleted, or added;
- (3) a guanosine triphosphate binding protein-coupled receptor protein encoded by a DNA which hybridizes with a DNA comprising the nucleotide sequence of SEQ ID NO: 2;
- (4) a guanosine triphosphate binding protein-coupled receptor protein encoded by a DNA which hybridizes with a DNA comprising the nucleotide sequence of SEQ ID NO: 21;
- (5) a partial peptide of the receptor protein as described in any one of (1) to (4);
- (6) a DNA encoding the receptor protein as described in any one of (1) to (4) or the partial peptide as described in (5);
- (7) the DNA described in (6), wherein said DNA comprising the nucleotide sequence of SEQ ID NO: 2 or NO: 21;
- (8) a vector containing the DNA as described in any one of (5) to (7);
- (9) a transformant carrying the vector as described in (8);
- (10) a method of producing the receptor protein as described in any one of (1) to (4) or the partial peptide as described in (5), the method comprising culturing the transformant as described in (9);
- (11) a method of screening for a ligand of the receptor protein as described in any one of (1) to (4), the method comprising exposing a test compound to the receptor protein as described in any one of (1) to (4) or the partial peptide as described in (5), and selecting a compound that binds to said protein or partial peptide;
- (12) a method of screening for a compound that inhibits the binding between the receptor protein as described in any one of (1) to (4) and its ligand, the method comprising,
  - (a) exposing a ligand to the receptor protein as described in any one of (1) to (4) or the partial peptide as described in (5) in the presence of a test compound, and detecting the binding activity between said protein or partial peptide and the ligand, and,
  - (b) comparing the binding activity detected in (a) with that in the absence of the test compound, and selecting a compound that reduces the binding activity between said protein or partial peptide and the ligand;
- (13) a kit for screening a compound that inhibits the binding between the receptor protein as described in any one of (1) to (4) and its ligand, the kit comprising the receptor protein as described in any one of (1) to (4) or the partial peptide as described in (5); and
- (14) an antibody which binds to the receptor protein as described in any one of (1) to (4).

**[0011]** "G protein-coupled receptor protein" herein refers to a receptor protein that transduces intracellular signals by activating G proteins. "Ligand" refers to a natural compound capable of binding to a G protein-coupled receptor and inducing signal transduction. "Agonist" refers to a compound having a bioactivity similar to that of the ligands of G protein-coupled receptors, including both natural and artificially synthesized compounds. "Antagonist" refers to a compound capable of inhibiting the bioactivity of a ligand of a G protein-coupled receptor, including both natural and artificially synthesized compounds. "Protein" and "peptide" as used herein include their salts as well.

**[0012]** The present invention relates to a novel G protein-coupled receptor protein. The nucleotide sequence of the cDNA encoding rat G protein-coupled receptor "BG2" isolated in the present invention is shown as SEQ ID NO: 2, and the amino acid sequence of the "BG2" protein is shown as SEQ ID NO: 1. The nucleotide sequence of the cDNA encoding human G protein-coupled receptor "BG2" isolated herein is shown as SEQ ID NO: 21, and the amino acid sequence of the human "BG2" protein is shown as SEQ ID NO: 20.

**[0013]** Rat "BG2" protein has 26%, 25%, and 29% homology to known G protein-coupled receptors; bovine muscarinic acetylcholine receptor M3 protein (Lee P.H. et al., Biochim. Biophys. Acta 1223:151-154 (1994)), human mus-

carinic acetylcholine receptor M5 protein (Bonner T.I. et al., Neuron 1:403-410 (1988)), and mouse  $\alpha$ 2A adrenoreceptor (Link R. et al., Mol. Pharmacol. 42:16-27 (1992)), respectively. The result of the hydrophobicity plot analysis showed that rat "BG2" protein contains hydrophobic regions (seven transmembrane regions) characteristic to G protein-coupled receptors. In addition, the coding region of the rat "BG2" cDNA has a size of approximately 1.2 kb, a size similar to that of the known G protein-coupled receptors.

**[0014]** Human "BG2" protein has 32%, 28%, and 27% homology to known G protein-coupled receptors; human  $\alpha$ -2C-1 adrenoreceptor (Regan J.W. et al., Proc. Natl. Acad. Sci. USA 85:6301-6305 (1988)), mouse  $\beta$ -1 adrenoreceptor (Jasper J.R. et al., Biochim. Biophys. Acta 1178:307-309 (1993)), and human muscarinic acetylcholine receptor M3 protein (Peralta E.G. et al., EMBO J 6:3923-3929 (1987)), respectively.

**[0015]** These results suggest that the "BG2" proteins belong to the G protein-coupled receptor family, which further suggests that they participate in signal transduction through the activation of G proteins upon ligand binding.

**[0016]** Furthermore, the result of a northern blot analysis showed that the genes encoding the "BG2" proteins are specifically expressed in the brain. *In situ* hybridization detected a strong expression of the rat "BG2" gene in the hippocampus and spinal cord, and also in the hypothalamus, thalamus, and cerebellum.

**[0017]** The hippocampus plays an important role in memory and learning, the cerebellum regulates the body motions, and the hypothalamus serves as the center of the autonomic nervous system. Thus, the "BG2" proteins are assumed to be involved in the regulation of these functions. Therefore, the "BG2" proteins and genes, or an agonist or antagonist that can regulate the "BG2" protein function(s) can be used in the treatment of disabilities in memory and learning, or the control of the autonomous nervous system, such as regulation of, blood pressure, digestion, body temperature, and food-intake etc.

**[0018]** "BG2" protein may be prepared as natural protein, and also as recombinant protein by using recombinant DNA technology. A natural protein may be prepared, for instance, by extracting tissues such as the thalamus and hypothalamus, speculated to express "BG2" protein, and performing immunoaffinity chromatography using anti-"BG2" antibody as described later on. On the other hand, the recombinant protein can be prepared by culturing transformant cells carrying DNA encoding "BG2" protein as described later on.

**[0019]** One skilled in the art can prepare an altered protein having a function (transduction of intracellular signals through G protein activation) equivalent to that of the natural protein by introducing modifications such as replacement of any amino acid contained in the natural rat or human "BG2" protein (SEQ ID NO: 1, or SEQ ID NO: 20, respectively) according to known methods. Mutations of amino acids in a protein may occur naturally. The G protein-coupled receptor protein of the present invention includes such mutant proteins having an amino acid sequence altered by replacement, deletion or addition, having a function equivalent to that of the natural protein. The methods of altering amino acids, known to one skilled in the art, include, the Kunkel method (Kunkel T.A. et al., Methods Enzymol. 154:367-382 (1987)), double primer method (Zoller M.J. and Smith M., Methods Enzymol. 154:329-350 (1987)), cassette mutation (Wells et al., Gene 34:315-323 (1985)), and megaprimer method (Sarkar G. and Sommer S.S., Biotechniques 8:404-407 (1990)). The number of mutated amino acids in a functionally equivalent protein is generally not more than 10% of all the amino acids, favorably not more than 10 amino acids, and more favorably not more than 3 amino acids (for instance, one amino acid).

**[0020]** One skilled in the art can also use hybridization techniques (Hanahan D. and Meselson M., Methods Enzymol. 100:333-342 (1983); Benton W.D. and Davis R.W., Science 196:180-182 (1977)) to isolate a highly homologous gene from various other species based on the sequence of the rat or human "BG2" cDNA (SEQ ID NO: 2, or SEQ ID NO: 21, respectively) or a part of it, and use the isolated DNA to obtain a protein a functionally equivalent to these "BG2" proteins. Thus, it is possible for one skilled in the art to prepare a protein functionally equivalent to the rat or human "BG2" proteins, which is encoded by a DNA that hybridizes to the rat or human "BG2" cDNA. The G protein-coupled receptor protein of the present invention includes these proteins as well. Species used for isolation of a functionally equivalent protein include mouse, rabbit, sheep, bovine, dog, and pig, for instance, and particularly the brain tissues such as thalamus and hypothalamus are suitable.

**[0021]** DNA encoding a functionally equivalent protein to the rat or human "BG2" proteins has usually high homology to the nucleotide sequence of the rat or human "BG2" cDNA (SEQ ID NO: 2, and NO: 21). High homology generally means a sequence identity of not less than 70%, favorably not less than 80%, and more favorably not less than 90% at the nucleotide level. The sequence identity can be determined by the FASTA program.

**[0022]** Hybridization to isolate a DNA having high homology to the rat or human "BG2" cDNA is usually performed in "6x SSC, 40% formamide, at 25°C" followed by a wash in "1x SSC at 55°C," favorably in "6x SSC, 40% formamide, at 37°C" followed by a wash in "0.2x SSC at 55°C," and more favorably in "6x SSC, 50% formamide, at 37°C" followed by a wash in "0.1x SSC at 62°C." One skilled in the art may easily obtain a hybridization condition similar to the above conditions by appropriately selecting factors such as the dilution rate of SSC, the formamide concentration, and temperature.

**[0023]** The present invention also includes a partial peptide of the above-described G protein-coupled receptor protein. The partial peptide of the present invention includes, for instance, those corresponding to the N-terminal region of

the G protein-coupled receptor protein, which can be utilized to prepare an antibody. The partial peptide of the invention has a length of at least 15 amino acids, and favorably 20 amino acids.

**[0024]** Furthermore, the present invention relates to a DNA encoding the G protein-coupled receptor protein of the invention as described above or its partial peptide. The DNA encoding the G protein-coupled receptor protein of the invention or its partial peptide includes cDNA, genomic DNA, and synthetic DNA, but is not limited as far as it encodes the protein or the peptide. cDNA encoding the G protein-coupled receptor of the invention can be obtained by screening a cDNA library from a tissue expressing the receptor (for instance, thalamus and hypothalamus) by using as a <sup>32</sup>P-labeled probe, the cDNA as described in SEQ ID NO: 2 or NO: 21, or a part of it, complementary RNA to the DNA, or a synthetic oligonucleotide comprising a part of the cDNA. Alternatively, cDNA may be cloned by synthesizing an oligonucleotide corresponding to the nucleotide sequence of the cDNA, and amplifying cDNA from an appropriate tissue (such as thalamus and hypothalamus) by PCR. Genomic DNA can be obtained by screening a genomic library by hybridization using as a <sup>32</sup>P-labeled probe, the cDNA as described in SEQ ID NO: 2 or NO: 21, or a part of it, complementary RNA to the DNA, or a synthetic oligonucleotide comprising a part of the cDNA. Alternatively, it may be cloned by synthesizing an oligonucleotide corresponding to the nucleotide sequence of the cDNA, and amplifying genome DNA by PCR. Synthetic DNA can be prepared by chemically synthesizing oligonucleotides comprising a part of the nucleotide sequence of SEQ ID NO: 2 or NO: 21, annealing them into a double strand, and ligating them using DNA ligase (Khorana H.G. et al., J. Biol. Chem. 251:565-570 (1976); Goeddel D.V. et al., Proc. Natl. Acad. Sci. USA 76:106-110 (1979)).

**[0025]** These DNA can be used for producing recombinant proteins. Namely, it is possible to prepare the G protein-coupled receptor protein of the invention as a recombinant protein by inserting a DNA encoding the receptor protein (DNA as described in SEQ ID NO: 2 or NO: 21, for instance) into an appropriate expression vector, culturing a transformant obtained by introducing the vector into an appropriate cell, and purifying the expressed protein. Since the G protein-coupled receptor protein of the invention is a receptor protein, it is possible to prepare it in a form expressed on the cell membrane.

**[0026]** Specifically, if the host is *Escherichia coli*, the plasmid vectors such as pET-3 (Rosenburg A.H. et al., Gene 56:125-135 (1987)) and pGEX-1 (Smith D.B. and Johnson K.S., Gene 67:31-40 (1988)) may be used. *E. coli* can be transformed by the Hanahan method (Hanahan D., J. Mol. Biol. 166:557-580 (1983)), electroporation (Dower W.J. et al., Nucleic Acids Res. 16:6127-6145 (1988)), and such. If the host is fission yeast (*Schizosaccharomyces pombe*), a plasmid vector such as pESP-1 (Lu Q. et al., Gene 200:135-144 (1997)) is used. Yeast can be transformed by spheroplast fusion (Beach D. and Nurse P., Nature 290:140 (1981)), and lithium acetate method (Okazaki K. et al., Nucleic Acids Res. 18:6485-6489 (1990)), etc.

**[0027]** If the host is a mammalian cell, such as Chinese Hamster ovary-derived CHO cells and human HeLa cells, vectors such as pMSG (Clontech) are used. Recombinant DNA is introduced into mammalian cells by calcium phosphate method (Graham F.L. and van der Eb A.J., J. Virology 52:456-467 (1973)), DEAE-dextran method (Sussman D.J. and Milman G., Mol. Cell. Biol. 4:1641-1643 (1984)), lipofection (Felgner P.L. et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987)), and electroporation (Neumann E. et al., EMBO J. 1:841-845 (1982)), etc. If the host is an insect cell, a baculovirus vector such as pBacPAK8/9 (Clontech) can be used. Transformation of insect cells is done by the methods described in the literature (BioTechnology 6:47-55 (1980)).

**[0028]** Recombinant proteins expressed in host cells can be purified by known methods. The proteins can also be synthesized as fusion proteins tagged with histidine residues at the N-terminus, or fused to glutathione-S-transferase (GST), and purified by using their binding ability toward a metal chelating resin, or a GST affinity resin (Smith M.C. et al., J. Biol. Chem. 263:7211-7215 (1988)), respectively. For instance, when the vector pESP-1 is used, the protein of interest is synthesized as a fusion protein with GST, which can be purified using GST affinity resin. The fusion protein may be digested with thrombin, or blood coagulating factor Xa to liberate the protein of interest.

**[0029]** Moreover, DNA encoding the G protein-coupled receptor protein of the present invention can be used in gene therapy of diseases that arise from a mutation of the protein. When used in gene therapy, the DNA can be introduced into human cells using retrovirus vectors (Danos O. and Mulligan R.C., Proc. Natl. Acad. Sci. USA 85:6460-6464 (1988); Dranoff et al., Proc. Natl. Acad. Sci. USA 90:3539-3543 (1993)), or adenovirus vectors (Wickham T.J. et al., Cell 73:309-319 (1993)), etc. To administer the vector to patients, transplantation of bone marrow, subcutaneous injection, and intravenous injection can be used (Asano S., Protein Nucleic acid and Enzyme 40:2491-2495 (1995)).

**[0030]** Furthermore, the present invention relates to an antibody that is capable of binding to the G protein-coupled receptor protein of the invention. Antibodies against the G protein-coupled receptor protein can be prepared by known methods in the art (for instance, refer to Shin-Seikagaku-Jikken-Kouza I: Protein I 389-406, Tokyo-Kagaku-Doujin). For instance, polyclonal antibodies are prepared as follows. An appropriate dose of the above protein or its partial peptide is administered into immune animals such as rabbits, guinea pigs, mice, or chickens. Administration may be performed together with the adjuvant (such as FIA or FCA) that promotes antibody production, and usually performed every couple of weeks. The titer of antibodies can be increased by performing multiple immunizations. After the final immunization, antisera are obtained by withdrawing blood from immune animals. Polyclonal antibodies are purified from antisera by

ammonium sulfate precipitation, fractionation by anion exchange chromatography, or affinity chromatography with either Protein A or immobilized antigen. Monoclonal antibodies are prepared as follows. The G protein-coupled receptor proteins of the invention or its partial peptide is administered into immune animals as described above. After the final immunization, their spleens or lymph nodes are excised. Then, antigen-producing cells are recovered from the spleens or the lymph nodes, and fused with myeloma cells using polyethylene glycol etc. to produce hybridomas. Desired hybridomas are selected by screening, and their culture supernatant is used to prepare monoclonal antibodies. Monoclonal antibodies can be purified by ammonium sulfate precipitation, fractionation by anion exchange chromatography, or affinity chromatography with either Protein A or immobilized antigen. Antibodies prepared thereby can be used not only in affinity purification of the G protein-coupled receptor protein of the invention, but also for the diagnosis or antibody treatment of diseases arising from the abnormal expression of the receptors, or detection of the expression level of the receptors.

[0031] If used for antibody treatment, humanized antibodies or human antibodies are preferable. Humanized antibodies, in case of mouse-human chimeric antibodies, are prepared by isolating the gene encoding the antibody against the G protein-coupled receptor protein from the producing mouse cells, replacing the constant region of the H chain of the antibody with that of the human IgE, and introducing it into mouse myeloma J558L cells (Neuberger M.S. et al., Nature 314:268-270 (1985)). Human antibodies can be prepared by immunizing mice, whose immune system is replaced with that of human, with the G protein-coupled receptor protein.

[0032] Furthermore, the present invention relates to a method of screening ligands of the G protein-coupled receptor protein of the invention. The method includes such processes as exposing a test compound to the G protein-coupled receptor protein or its partial peptide, and selecting compounds that are capable of binding to the proteins or the peptide. Compounds to be tested include known compounds such as acetylcholine, adenosine, adrenaline, noradrenaline, angiotensin, bombesin, bradykinin, C5a, anaphylatoxin, calcitonin, cannabinoids, chemokines, cholecystokinin, dopamine, endothelin, formylmethionylpeptide, GABA, galanin, glucagon, glutamate, glycopeptide hormone, histamine, 5-hydroxytryptophan, leucotriene, melanocortin, neuropeptide Y, neurotensin, odorant, opioid peptide, opsin, parathyroid hormone, platelet activating factor, prostanoid, somatostatin, tachykinin, thrombin, thyrotropin releasing hormone, vasopressin, and oxytocin (Watson S. and Arkinstall S., G protein Linked Receptor FactsBook, Academic Press (1994)), and also other purified proteins, expressed products of genes (including libraries), extracts of tissues or cells in which the ligand is stipulated to be expressed (the brain, thalamus, and hypothalamus etc.), and the culture medium of the cells. The G protein-coupled receptor proteins may be used in a form expressed in desired cells (including transformants genetically engineered to express the proteins) or on the cell surface, in a form of the membrane fractions of the cells, or in a form bound to an affinity column. If necessary, test compounds may be labeled appropriately. Methods for labeling include radioisotope labeling, and fluorescence labeling, but not limited thereto. The binding between the G protein-coupled receptor proteins and test compounds can be examined by detecting the label added to the compound (for instance, measuring the radioactivity or fluorescence intensity), or using as an index, intracellular signaling triggered by the compound binding to the G protein-coupled receptor protein (such as G protein activation, the change in the concentration of  $Ca^{2+}$  or cAMP, phospholipase C activation, and the change in pH). Specific methods can be employed as described in the literatures (Cell Calcium 14:663-671 (1993); Analytical Biochemistry 226:349-354 (1995); J. Biol. Chem. 268:5957-5964 (1993); Cell 92:573-585 (1998); Nature 393:272-273 (1998)), and unexamined published Japanese patent application (JP-A) No. Hei 9-268). Alternatively, the binding may be detected by measuring the activity of the reporter gene using two-hybrid system (Zervos et al., Cell 72:223-232 (1994); Fritz et al., Nature 376:530-533 (1995)).

[0033] The present invention also relates to a method of screening for a compound which can inhibit the binding between the G protein-coupled receptor proteins of the invention and their ligands. The method includes processes of (a) exposing the ligand to the G protein-coupled receptors or their partial peptides in the presence of test compound, and detecting the binding ability between the ligand and the proteins or peptides, and (b) comparing the ability detected in (a) with that in the absence of the compound, and selecting compounds which are capable of decreasing the binding ability. Compounds to be tested include proteins, peptides, non-peptide compounds, artificially synthesized compounds, extracts of tissues and cells, sera, but not limited thereto. The G protein-coupled receptor proteins may be used in a form expressed in desired cells (including transformants genetically engineered to express the proteins) or on the cell surface, in a form of the membrane fractions of the cells, or in a form bound to an affinity column. If necessary, tested compounds may be labeled appropriately. Methods for labeling include radioisotope labeling, and fluorescence labeling, but not limited thereto. The binding between the G protein-coupled receptor proteins and test compounds can be examined by detecting the label added to the compound (for instance, measuring the radioactivity or fluorescence intensity), or using intracellular signaling, as an index, that are triggered by the compound binding to the G protein-coupled receptor protein (such as G protein activation, the change in the concentration of  $Ca^{2+}$  or cAMP, phospholipase C activation, and the change in pH). Specific methods can be employed as described in the literatures (Cell Calcium 14:663-671 (1993); Analytical Biochemistry 226:349-354 (1995); J. Biol. Chem. 268:5957-5964 (1993); Cell 92:573-585 (1998); Nature 393:272-273 (1998)), and JP-A Hei 9-268). If the result of the detection showed that the binding

activity in the presence of a test compound is lower than that in the absence of the compound, the compound is judged to be capable of inhibiting the binding between the G protein-coupled receptors or their partial peptides and their ligands. These compounds include those capable of triggering the intracellular signaling through binding to the G protein-coupled receptor (agonist), and those not having such activity (antagonist). Agonists have similar bioactivities to those of the ligands of the G protein-coupled receptors. On the other hand, antagonists inhibit the bioactivities of the ligands. Therefore, these agonists and antagonists are useful as medicinal components for treatment of diseases arising from disorders in the signaling pathway mediated by the G protein-coupled receptors.

[0034] Furthermore, the present invention relates to a screening kit for compounds that inhibit the binding between the G protein-coupled receptor proteins and their ligands. The G protein-coupled receptor proteins or their partial peptides may be in a form expressed in desired cells (including transformants genetically engineered to express the protein) or on the cell surface, in a form of membrane fractions of the cell, or in a form bound to an affinity column. Components of the kit of the invention may include, other than the above described receptor protein samples, ligand samples (both labeled and unlabeled), and buffers for the reaction between the ligand and the receptor protein, and wash solutions. Labels to be added to the ligands include radioisotope and fluorescence, for instance. The kit of the invention can be used as described in JP-A Hei 9-268.

#### Brief Description of the Drawings

#### [0035]

Figure 1 shows the hydrophobicity plot of the mouse BG2 protein. The seven hydrophobic regions (transmembrane regions) that are characteristics of the G protein-coupled receptor proteins are indicated by the numbers from 1 to 7. The numbers in the bottom indicate those of the amino acid residues in the "BG2" protein.

Figure 2 shows the result of northern blot analysis of the tissue specific expression of the human and mouse BG2 genes.

Figure 3 shows the result of *in situ* hybridization analysis of the location of the mouse "BG2" gene expression in the brain.

Figure 4 shows the result of *in situ* hybridization analysis of the location of the mouse "BG2" gene expression in the spinal cord. "Sense" and "Antisense" indicate the results using sense RNA probe (not hybridizing with mRNA; negative control), and antisense RNA probe (hybridizing with mRNA), respectively.

#### Best Mode for Carrying out the Invention

[0036] The present invention is illustrated in detail below with reference to examples, but is not to be construed as being limited thereto.

#### Example 1: Isolation of a gene encoding a rat G protein-coupled receptor

[0037] The G protein-coupled receptors share a characteristic structure composed of seven transmembrane regions, and the amino acid sequences of the transmembrane regions and the adjacent regions are well conserved. The present inventors first compared the nucleotide sequences of the second and the seventh transmembrane domains, which are highly conserved, with known G protein-coupled receptors: mouse neuropeptide Y receptor Y1 (GenBank Accession Number Z18280), rat Y1 (Z11504), human Y1 (M84755), mouse neuropeptide Y receptor Y4 (U40189), rat Y4 (Z68180), human Y4 (Z66526), and mouse neuropeptide Y receptor Y6 (U58367), and synthesized novel sense and antisense primers, as described in SEQ ID NO: 3 and NO: 4, respectively.

[0038] Next, single stranded cDNA was synthesized from poly(A)<sup>+</sup>RNA prepared from rat thalamus and hypothalamus using the RNA-PCR kit (TaKaRa), and PCR was performed using the two primers. Specifically, poly(A)<sup>+</sup>RNA was prepared from rat thalamus and hypothalamus using Fasttrack 2.0 kit (Invitrogen). Then, 75 ng of the poly(A)<sup>+</sup>RNA was used to synthesize complementary DNA according to the protocol accompanying the RNA-PCR kit (TaKaRa). PCR amplification was performed using all the cDNA. The reaction mixture comprising each 0.15 mM dNTPs, 1.5 mM MgCl<sub>2</sub>, 0.025 U/μl rTaq polymerase (TaKaRa), each 0.5 μM degenerated primer Fg (SEQ ID NO: 3) and Rb (SEQ ID NO: 4), and enzyme accompanying 10x PCR buffer was prepared making a total of 130 μl, and aliquoted into six 20 μl fractions. PCR was performed with the Pertier thermal cycler PTC200 (MJ Research) under conditions as follows: a single cycle of 94°C for 2 min, followed by 35 cycles of 94°C for 30 sec, 48°C for 1 min, 72°C for 1 min 30 sec, and then a single cycle of 72°C for 8 min. After PCR, the six reaction-solutions were combined, and the amplified products were purified using the Wizard PCR purification kit (Promega), and then eluted with 30 μl TE. 2 μl of the TE eluate was used for cloning into the pCR2.1 vector of the TOPO TA cloning kit (Invitrogen). XL1-Blue cells were used as the host cell and transformed using the *E. coli* pulser (BioRad). From the resulting transformants, 5,760 colonies having white or light blue

color were randomly selected using the gene library construction system BioPick (BioRobotics), and inoculated into fifteen 384-well plates containing LB media supplemented with 100 µg/ml of ampicillin. Clones were cultured at 37°C overnight, and replica plated onto a filter on top of a LB agar plate containing 100 µg/ml ampicillin and 25% glycerol, and another filter on top of a LB agar plate containing 100 µg/ml ampicillin, for preparing a glycerol stock, and colony hybridization, respectively, using a gene library replicating system BioGrid (BioRobotics).

**[0039]** Since the obtained PCR clones were expected to contain multiple overlapping clones of the NPY receptor cDNA, 80 clones out of the 5,760 clones were randomly selected, and their nucleotide sequences were partially determined. To determine the nucleotide sequence, plasmid DNA purified by the plasmid automatic isolating system PI100sigma (Kurabo) was used as a template. The sequence reactions were performed using the dye-primer-cycle sequencing kit FS (Perkin Elmer), and the reaction products were separated by electrophoresis using the DNA sequencer 377 (Perkin Elmer). The homology search of the obtained sequence using the blast program of the Wisconsin package (Genetic Computer Group) showed that 29 out of the 80 clones were the cDNAs encoding the coiled-coil like protein 1 (GeneBank Accession Number U79024) while 17 clones were those of the neuropeptide Y receptor Y1 (Z11504). Then, these two cDNA fragments were used as a probe for hybridization with the filters containing a library of the degenerated PCR amplified fragments. Probes were prepared by amplifying the insert of the respective clones by PCR, purifying the products using the Wizard PCR purification kit (Promega), and labeling them with [ $\alpha$ -<sup>32</sup>P] dCTP using the Prime-It II random primer labeling kit (Stratagene). Colony hybridization was performed according to the standard method (Sambrook et al., Molecular Cloning: A laboratory manual 2<sup>nd</sup> edition (1989)). Colonies that were negative for either the coiled-coil like protein 1 or the neuropeptide Y receptor Y1 were selected and their partial nucleotide sequences were determined. For DNA sequencing, the insert fragment of each clone was amplified by PCR from the culture medium, purified using the PCR product purification kit (Amersham), and used as a template. The sequence reactions were performed using the dye-primer-cycle sequencing kit FS (Perkin Elmer), and the reaction products were separated by electrophoresis using the DNA sequencer 377 (Perkin Elmer). The obtained sequences were analyzed by the homology search using the blast program of the Wisconsin package (Genetic Computer Group), and, as a result, a clone which has significant homology to the muscarinic acetylcholine receptor M5 (GeneBank Accession Number M22926) was found. The clone has been deposited at the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology.

Name of the depositary institution: National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, MITI.

Address of the depositary institution: 1-1-3 Higashi, Tsukuba, Ibaraki 305-8566, Japan.

Date of deposit: 12/25/1997.

Accession Number: FERM BP-6575

**[0040]** Next, in order to isolate the full-length cDNA of the gene, cDNA libraries were prepared from rat thalamus and hypothalamus. cDNA was synthesized according to the protocol accompanying the cDNA synthesizing kit (Stratagene), and the vector pEF1x and the host XL1-blue MRF<sup>+</sup> (Stratagene) were used. The pEF1x is a refined derivative of the pcDNA3 (Invitrogen) prepared as follows.

(1) Preparation of the human EF1a promoter (GeneBank Accession Number J04617)

**[0041]** PCR was performed using human genomic DNA with primers (SEQ ID NO: 6/ CGAGGATCCGTGAGGCTC-CGGTGCCCGTC; SEQ ID NO: 7/ CGGGTAAGCTTCACGACACCTGAAATGGAAGA). The products were digested with BamHI (TaKaRa) and HindIII (TaKaRa), and subcloned into the plasmid vector pUC19 (TaKaRa). The resulting plasmid was digested with XhoI, blunt-ended with the Klenow fragment (TaKaRa), and self-ligated using the DNA ligation kit (TaKaRa). The resulting plasmid was digested with BamHI and HindIII, and the insert was recovered.

(2) Alteration of the pcDNA3

**[0042]** The pcDNA3 was digested with MluI (TaKaRa), blunt-ended with the Klenow fragment (TaKaRa), and self-ligated using the DNA ligation kit. The resulting plasmid was digested with AflIII (New England Biolabs) and SmaI (TaKaRa), blunt-ended with the Klenow fragment (TaKaRa), and self-ligated using the DNA ligation kit. Then, the obtained plasmid was digested with BglII (TaKaRa) and HindIII, and the fragment depleted of the CMV promoter was recovered, and ligated with the insert fragment as described in (1) using the DNA ligation kit to construct the pEF1x.

**[0043]** Next, the nucleotide sequence of the cDNA fragment was used to synthesize oligonucleotide probe (SEQ ID NO: 8/ CCTTCTGCATCCCATTTGTACGTACC), and multiple clones were obtained from the above cDNA libraries from rat thalamus and hypothalamus according to the protocol of the gene trapper cDNA positive selection system (GIBCO BRL). Then, colony hybridization was performed using the cDNA insert of the above isolated clone (FERM BP-6574)



as a probe, and a positive clone was obtained. This clone has been deposited at the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology.

Name of the depositary institution: National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, MITI.

Address of the depositary institution: 1-1-3 Higashi, Tsukuba, Ibaraki 305-8566, Japan.

Date of deposit: 12/25/1997.

Accession Number: FERM BP-6574

**[0044]** The insert fragment of the clone was of 2.7 kb. Plasmid DNA was prepared by the QIAprep Midi Kit (QIAGEN), and the complete nucleotide sequence was determined using the shotgun cloning method (Sambrook et al., Molecular Cloning: A laboratory manual 2<sup>nd</sup> edition (1989)). cDNA fragmentation was performed using the closed sonifier biomaterial treating system Biorupter (Tousou Denki), and the DNA fragments were separated by electrophoresis on a 2% agarose gel. Fragments of around 0.6 kb were purified using the gene clean spin kit (bio101), blunt-ended with T4 DNA polymerase (TaKaRa), and cloned into the HincII-BAP treated pUC118 vector, XL1-Blue was used as a host cell, and transformed using the *E.coli* pulser (BioRad). The obtained shotgun clones were sequenced using the dye-primer cycle sequencing kit FS (Perkin Elmer), or the dye-terminator cycle sequencing kit (Perkin Elmer). The resulting sequences were combined and edited to get the complete nucleotide sequence using the DNA sequencing software Sequencher (Hitachi Software). The complete nucleotide sequence is composed of 2700 bp, and turned out to be encoding a protein of 413 amino acids (SEQ ID NO: 5). Because there is a stop codon in the 5' region of the open reading frame, the cDNA is supposed to include the entire coding region (SEQ ID NO: 2). When this sequence was translated into the amino acid sequence, the hydrophobicity plot identified seven transmembrane regions from 1 to 7 (Figure 1).

**[0045]** In addition, the open reading frame size was approximately 1.2 kb, which is similar to that of the known G protein-coupled receptors. G protein-coupled receptor proteins have common features in their amino acid sequences, and thus form a protein family. As a result of the homology search using the amino acid sequence encoded by the isolated cDNA, the encoded protein was found to be a novel receptor protein having a homology of 26%, 25%, and 29% to known G protein-coupled receptors: bovine muscarinic acetylcholine receptor M3 protein (Lee P.H. et al., Biochim. Biophys. Acta 1223:151-154 (1994)), human muscarinic acetylcholine receptor M5 protein (Bonner T.I. et al., Neuron 1:403-410 (1988)), and mouse  $\alpha 2A$  adrenoceptor (Link R. et al., Mol. Pharmacol. 42:16-27 (1992)), respectively.

#### Example 2: The isolation of the human G protein-coupled receptor gene

**[0046]** The obtained rat sequence was subjected to EST search to reveal a fragment of the human homologue (gene bank NID: 946030 and NID: 901756). Human fetal brain cDNA was amplified by PCR using the specific primers IF01 (SEQ ID NO: 9/ CTTCCGCCGGCCTTCACCAA) and IR02 (SEQ ID NO: 10/ ACAGACACGGCGGGGCTCAC) (probe 1). A human  $\lambda$  EMBL3 SP6/T7 genomic library (Clontech) of a size of  $1.2 \times 10^6$  pfu was screened using probe 1 according to standard procedure plaque hybridization procedures. Two positive clones were thus isolated. The obtained phage-clones were digested with SacI, and three bands of a clone were subcloned. These fragments, termed I1 (SEQ ID No: 11), I3 (SEQ ID No: 12) and I5 (SEQ ID No: 13), were sequenced and a hypothetical sequence was speculated by comparing with the rat homologue. I1 and I3 were subjected to PCR amplification using specific primers YS03 (SEQ ID NO: 14/ TGAACGCTTCGGGGGCGCTG) and YS05 (SEQ ID NO: 15/ GAGATGGCGAGGTTGAGCAGG), YS12 (SEQ ID NO: 16/ GGCTCCAAGCCATCGGCGTC) and YS14 (SEQ ID NO: 17/ CTCACCTCCAGCAGTGCTCC) and the PCR products were termed probe 2 and probe 3, respectively. Human hypothalamus cDNA ( $1.3 \times 10^6$  phage) was plated at a density of  $5.6 \times 10^4$  pfu/150mm plate. The obtained sub-pools were checked by PCR using the primers YS03 and YS05. One positive sub-clone was screened in the same method as the screening of the genomic library, using probe 2. One cDNA clone containing 5'UTR to TM5 was obtained and named cDNA clone 1.

**[0047]** Probe 4 was amplified by PCR from cDNA clone 1 using the primers YS07 (SEQ ID NO: 18/ GCCTCCG-CACCCAGAACAAC) and YS10 (SEQ ID NO: 19/ TGCGCCTCTGGATGTTTCA). Phase screening of the human hippocampus library ( $3 \times 10^6$  pfu) was done in the same method as the genomic library, using probe 3 and probe 4. A few clones were obtained and the longest one, termed cDNA clone 2, was sequenced. It has the region between TM2 and 3'UTR. cDNA clone 1 was digested by SacII, and the 3.3kb band, which contained vector and the 5'-end region, was treated by shrimp alkaline phosphatase. cDNA clone 2 was also digested by SacII, and the 1.7kb fragment was ligated into the 3.3kb fragment from cDNA clone 1. The clone into which this ligated fragment was inserted has been deposited at the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology.

Name of the depositary institution: National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, MITI.

Address of the depositary institution: 1-1-3 Higashi, Tsukuba, Ibaraki 305-8566, Japan.

Date of deposit: 12/17/1998.

Accession Number: FERM BP-6609

5 [0048] Human "BG2" cDNA nucleotide sequence is shown in SEQ ID NO: 21, and the amino acid sequence of the protein encoded by the said cDNA in SEQ ID NO: 20.

[0049] Human "BG2" protein had 32%, 28%, and 27% homology to known G protein-coupled receptors; human  $\alpha$ -2C-1 adrenoreceptor (Regan J.W. et al., Proc. Natl. Acad. Sci. USA 85:6301-6305 (1988)), mouse  $\beta$ -1 adrenoreceptor (Jasper J.R. et al., Biochim. Biophys. Acta 1178:307-309 (1993)), and human muscarinic acetylcholine receptor M3 protein (Peralta E.G. et al., EMBO J. 6:3923-3929 (1987)), respectively.

#### Example 3: Northern blot analysis

[0050] Probe 4 was labeled with  $^{32}$ P $\gamma$ -dCTP (Amersham, Prime It II) and used as cDNA probe for the detection of human "BG2". Human Multiple Tissue Northern (MTN) Blots Membrane with Express HybriTM was purchased from Clontech. After prehybridization of the membrane at 68°C for 30 min, it was hybridized with the probe at 68°C for 1 hr (final concentration of the probe was  $1.5 \times 10^6$  cpm/ml.) The blot was rinsed with 2 x SSC containing 0.1% SDS at 42°C for 30 min, and the final wash was done at 50°C for 30 min in 0.1 x SSC containing 0.1% SDS. The blot was then exposed at -80°C for 2.5 days to Kodak autoradiographic film.

20 [0051] For the detection of the mouse "BG2," probe was prepared by PCR amplifying using the rat "BG2" cDNA as a template with sense primer MF2 (SEQ ID NO: 22/ TGCATCCCATTGTACGTNCC), and antisense primer MR1 (SEQ ID NO: 24/ TGCTCTGGGACACCATCTTC), purifying the amplified products by electrophoresis on an agarose gel, and labeling them as above described for human gene.

[0052] Blotting membrane used was the Rat MTN (Multiple Tissue Northern) blot (Clontech). Hybridization was performed at 42°C overnight in hybridization buffer (50% formamide, 4x SSPE, 1% SDS, 0.5% BLOTTO, and 100  $\mu$ g/ml salmon sperm DNA). The membrane was washed at 65°C in 0.1x SSC containing 0.1% SDS, and then exposed to the Kodak autoradiography film at -80°C overnight.

[0053] The result showed that the human and rat "BG2" genes are strongly expressed particularly in the brain (Fig. 2).

#### Example 4: *In situ* Hybridization

[0054] Adult male Sprague-Dawley rats (Charles River Japan) aged 13-18 weeks were anesthetized with inhalation of ether, connected to a rotary pump and infused with chilled 4% paraformaldehyde in phosphate buffer (pH7.2) via a cannula inserted into the left ventricle. After perfusion, brain, pituitary and spinal cord were removed and dissected to sagittal or coronal sections. The tissue specimens were postfixed with the same fixative overnight at 4°C. The following process was carefully done to avoid RNA contamination. Tissue specimens were embedded in paraffin wax in a routine manner, then paraffin sections were cut into a thickness of 6  $\mu$ m by rotary microtome (Model HM 355; MICROM Laborgerate GmbH). The sections were stored in moisture free condition at -20°C until proceeded to *in situ* hybridization.

40 [0055] For preparation of rat BG2 sense and antisense RNA probes, the cDNA fragment, amplified by PCR from MP-21 plasmid DNA using a sense primer MF2 (SEQ ID NO:22/ TGCATCCCATTGTACGTNCC) and antisense primer MR3 (SEQ ID NO:23/ ATCATTAGGAGCGTGTANGG), was cloned into pZErO-2 vector (Invitrogen). The RNA probes were labeled with digoxigenin using DIG RNA Labeling Kit (Boehringer Mannheim). Paraffin sections were de-paraffinized with xylene and transferred to distilled water after rinsing with graded ethyl alcohol. *In situ* Hybridization Reagents (ISHR, Code No. 316-01951; Nippon Gene) were used as reagents without digoxigenin-labeled RNA. The sections were incubated with two changes of phosphate buffer saline (PBS; ISHR 1) for 1 min and 10 min. The sections were treated with proteinase K (ISHR 6) for 10 min at 37°C. Acetylation was done with acetylation buffer (ISHR 3) containing acetic anhydride (ISHR 4) for 15 min, followed by quenching with PBS/glycine buffer (ISHR 2) for 20 min at room temperature, rinsed twice with 4 x SSC (ISHR 5) for 10 min and then rinsed with PBS buffer for 10 min. After pre-hybridization with 50% formamide/2 x SSC for 30 min at room temperature, hybridization was performed for 16 hr at 42 °C using digoxigenin-labeled RNA probe (1  $\mu$ g/ml).

50 [0056] Post hybridization washing was performed twice 50% formamide/2 x SSC for 10 min at 42°C. Then the sections were treated with RNase A (ISHR10)/NET buffer (ISHR 9) for 30 min at 37°C after rinsing with NET buffer (ISHR 9) for 5 min at 37°C. After washing twice with 0.1 x SSC buffer (ISHR 11) for 20 min, the sections were transferred and labeled digoxigenin was detected using the Digoxigenin Detection Kit (Boehringer Mannheim). Then the sections were rinsed with 100 mM Tris-HCl, 150 mM NaCl containing buffer (Buffer 1) for 1 min at room temperature then incubated with blocking reagent (Buffer 2) for 30 min at room temperature. The sections were incubated with anti-digoxigenin alka-

line phosphatase-labeled antibody for 60 min at room temperature. After washing with Buffer 1 for 15 min and Buffer 3 for 2 min at room temperature, the solutions were incubated with NBT/X-phosphate solution diluted with Buffer 3 for 12-14 hr at room temperature. The sections were mounted with glycerol or Permount after washing with Buffer 4.

5 [0057] As a result, as shown in Figures 3 and 4, BG2 cDNA probe was strongly hybridized to the hippocampus and the spinal cord. Hybridization of a medium extent was also detected in hypothalamus and cerebellum.

Industrial Applicability

10 [0058] The present invention has provided novel G protein-coupled receptor proteins specifically expressed in the brain, and their genes. Use of the receptors makes it possible to screen their ligands and compounds that are candidates for medicines. These ligands and candidate compounds would be useful in the diagnosis and treatment of diseases arising from disorders of signal transduction pathway mediated by the G protein-coupled receptor of the invention.

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

5

<110> BANYU PHARMACEUTICAL CO., LTD.

10

<120> GUANOSINE TRIPHOSPHATE (GTP) BINDING PROTEIN-COUPLED RECEPTOR  
PROTEIN

<130> B1-902PCT

15

<140>

<141>

<150> JP 1997-361187

20

<151> 1997-12-26

<160> 24

25

<170> PatentIn Ver. 2.0

<210> 1

<211> 413

30

<212> PRT

<213> Rattus norvegicus

<400> 1

35

Met Glu Arg Ala Pro Pro Asp Gly Leu Met Asn Ala Ser Gly Thr Leu  
1 5 10 15

40

Ala Gly Glu Ala Ala Ala Ala Gly Gly Ala Arg Gly Phe Ser Ala Ala  
20 25 30

45

Trp Thr Ala Val Leu Ala Ala Leu Met Ala Leu Leu Ile Val Ala Thr  
35 40 45

Val Leu Gly Asn Ala Leu Val Met Leu Ala Phe Val Ala Asp Ser Ser  
50 55 60

50

Leu Arg Thr Gln Asn Asn Phe Phe Leu Leu Asn Leu Ala Ile Ser Asp  
65 70 75 80

55

EP 1 043 395 A1

Phe Leu Val Gly Ala Phe Cys Ile Pro Leu Tyr Val Pro Tyr Val Leu  
 85 90 95  
 5  
 Thr Gly Arg Trp Thr Phe Gly Arg Gly Leu Cys Lys Leu Trp Leu Val  
 100 105 110  
 10  
 Val Asp Tyr Leu Leu Cys Ala Ser Ser Val Phe Asn Ile Val Leu Ile  
 115 120 125  
 Ser Tyr Asp Arg Phe Leu Ser Val Thr Arg Ala Val Ser Tyr Arg Ala  
 15 130 135 140  
 Gln Gln Gly Asp Thr Arg Arg Ala Val Arg Lys Met Ala Leu Val Trp  
 145 150 155 160  
 20  
 Val Leu Ala Phe Leu Leu Tyr Gly Pro Ala Ile Leu Ser Trp Glu Tyr  
 165 170 175  
 25  
 Leu Ser Gly Gly Ser Ser Ile Pro Glu Gly His Cys Tyr Ala Glu Phe  
 180 185 190  
 Phe Tyr Asn Trp Tyr Phe Leu Ile Thr Ala Ser Thr Leu Glu Phe Phe  
 30 195 200 205  
 Thr Pro Phe Leu Ser Val Thr Phe Phe Asn Leu Ser Ile Tyr Leu Asn  
 210 215 220  
 35  
 Ile Gln Arg Arg Thr Arg Leu Arg Leu Asp Gly Gly Arg Glu Ala Gly  
 225 230 235 240  
 Pro Glu Pro Pro Pro Asp Ala Gln Pro Ser Pro Pro Pro Ala Pro Pro  
 40 245 250 255  
 Ser Cys Trp Gly Cys Trp Pro Lys Gly His Gly Glu Ala Met Pro Leu  
 45 260 265 270  
 His Ser Ser Gly Ser Ser Ser Arg Gly Thr Glu Arg Pro Arg Ser Leu  
 275 280 285  
 50  
 Lys Arg Gly Ser Lys Pro Ser Ala Ser Ser Ala Ser Leu Glu Lys Arg  
 290 295 300  
 55

EP 1 043 395 A1

5 Met Lys Met Val Ser Gln Ser Ile Thr Gln Arg Phe Arg Leu Ser Arg  
 305 310 315 320  
 Asp Lys Lys Val Ala Lys Ser Leu Ala Ile Ile Val Ser Ile Phe Gly  
 325 330 335  
 10 Leu Cys Trp Ala Pro Tyr Thr Leu Leu Met Ile Ile Arg Ala Ala Cys  
 340 345 350  
 15 His Gly Arg Cys Ile Pro Asp Tyr Trp Tyr Glu Thr Ser Phe Trp Leu  
 355 360 365  
 Leu Trp Ala Asn Ser Ala Val Asn Pro Val Leu Tyr Pro Leu Cys His  
 370 375 380  
 20 Tyr Ser Phe Arg Arg Ala Phe Thr Lys Leu Leu Cys Pro Gln Lys Leu  
 385 390 395 400  
 25 Lys Val Gln Pro His Gly Ser Leu Glu Gln Cys Trp Lys  
 405 410  
 30 <210> 2  
 <211> 1239  
 <212> DNA  
 35 <213> Rattus norvegicus  
 <220>  
 <221> CDS  
 40 <222> (1)..(1239)  
 <400> 2  
 atg gag cgc gcg ccg ccc gac ggg ctg atg aac gcg tcg ggc act ctg 48  
 45 Met Glu Arg Ala Pro Pro Asp Gly Leu Met Asn Ala Ser Gly Thr Leu  
 1 5 10 15  
 gcc gga gag gcg gcg gct gca ggc ggg gcg cgc ggc ttc tcg gct gcc 96  
 50 Ala Gly Glu Ala Ala Ala Ala Gly Gly Ala Arg Gly Phe Ser Ala Ala  
 20 25 30  
 55

EP 1 043 395 A1

	tgg acc gct gtc ctg gct gcg ctc atg gcg ctg ctc atc gtg gcc aca	144
5	Trp Thr Ala Val Leu Ala Ala Leu Met Ala Leu Leu Ile Val Ala Thr	
	35 40 45	
	gta ctg ggc aac gcg ctg gtc atg ctc gcc ttc gtg gcg gat tgg agc	192
10	Val Leu Gly Asn Ala Leu Val Met Leu Ala Phe Val Ala Asp Ser Ser	
	50 55 60	
	ctc cgc acc cag aac aac ttc ttt ctg ctc aac ctc gcc atc tcc gac	240
15	Leu Arg Thr Gln Asn Asn Phe Phe Leu Leu Asn Leu Ala Ile Ser Asp	
	65 70 75 80	
	ttc ctc gtg ggt gcc ttc tgc atc cca ttg tac gta ccc tat gtg ctg	288
20	Phe Leu Val Gly Ala Phe Cys Ile Pro Leu Tyr Val Pro Tyr Val Leu	
	85 90 95	
	acc gcc cgt tgg acc ttc gcc cgg gcc ctc tgc aag ctg tgg ctg gtg	336
25	Thr Gly Arg Trp Thr Phe Gly Arg Gly Leu Cys Lys Leu Trp Leu Val	
	100 105 110	
	gta gac tac cta ctg tgt gcc tcc tgg gtc ttc aac atc gta ctc atc	384
30	Val Asp Tyr Leu Leu Cys Ala Ser Ser Val Phe Asn Ile Val Leu Ile	
	115 120 125	
	agc tat gac cga ttc ctg tca gtc act cga gct gtc tcc tac agg gcc	432
35	Ser Tyr Asp Arg Phe Leu Ser Val Thr Arg Ala Val Ser Tyr Arg Ala	
	130 135 140	
	cag cag ggg gac acg aga cgg gcc gtt cgg aag atg gca ctg gtg tgg	480
40	Gln Gln Gly Asp Thr Arg Arg Ala Val Arg Lys Met Ala Leu Val Trp	
	145 150 155 160	
	gtg ctg gcc ttc ctg ctg tat ggg cct gcc atc ctg agt tgg gag tac	528
45	Val Leu Ala Phe Leu Leu Tyr Gly Pro Ala Ile Leu Ser Trp Glu Tyr	
	165 170 175	
	ctg tct ggt gcc agt tcc atc ccc gag gcc cac tgc tat gct gag ttc	576
50	Leu Ser Gly Gly Ser Ser Ile Pro Glu Gly His Cys Tyr Ala Glu Phe	
	180 185 190	
	ttc tac aac tgg tac ttt ctc atc acg gcc tcc acc ctc gag ttc ttc	624

55

EP 1 043 395 A1

Phe Tyr Asn Trp Tyr Phe Leu Ile Thr Ala Ser Thr Leu Glu Phe Phe  
195 200 205

5  
acg ccc ttc ctc agc gtt acc ttc ttc aac ctc agc atc tac ctg aac 672  
Thr Pro Phe Leu Ser Val Thr Phe Phe Asn Leu Ser Ile Tyr Leu Asn  
210 215 220

10  
atc cag agg cgc acc cgc ctt cgg ctt gat ggg ggc cgt gag gct ggc 720  
Ile Gln Arg Arg Thr Arg Leu Arg Leu Asp Gly Gly Arg Glu Ala Gly  
225 230 235 240

15  
cca gaa ccc cca cca gat gcc cag ccc tcg cca cct cca gct ccc ccc 768  
Pro Glu Pro Pro Pro Asp Ala Gln Pro Ser Pro Pro Pro Ala Pro Pro  
245 250 255

20  
agc tgc tgg ggc tgc tgg cca aaa ggg cat ggc gag gcc atg ccg ttg 816  
Ser Cys Trp Gly Cys Trp Pro Lys Gly His Gly Glu Ala Met Pro Leu  
260 265 270

25  
cac agc tct ggc agc tcc tca agg ggc act gag agg cca cgc tca ctc 864  
His Ser Ser Gly Ser Ser Ser Arg Gly Thr Glu Arg Pro Arg Ser Leu  
275 280 285

30  
aaa agg ggc tcc aag cca tca gca tct tca gca tcc ctg gag aag cgc 912  
Lys Arg Gly Ser Lys Pro Ser Ala Ser Ser Ala Ser Leu Glu Lys Arg  
290 295 300

35  
atg aag atg gtg tcc cag agc atc acc cag cgc ttc cgg ctg tcg cgg 960  
Met Lys Met Val Ser Gln Ser Ile Thr Gln Arg Phe Arg Leu Ser Arg  
305 310 315 320

40  
gac aag aag gtg gcc aag tcg ctg gcc atc atc gtg agc atc ttt ggg 1008  
Asp Lys Lys Val Ala Lys Ser Leu Ala Ile Ile Val Ser Ile Phe Gly  
325 330 335

45  
ctc tgc tgg ggc ccg tac acg ctc cta atg atc atc cga gct gct tgc 1056  
Leu Cys Trp Ala Pro Tyr Thr Leu Leu Met Ile Ile Arg Ala Ala Cys  
340 345 350

50  
cat ggc cgc tgc atc ccc gat tac tgg tac gag acg tcc ttc tgg ctt 1104  
His Gly Arg Cys Ile Pro Asp Tyr Trp Tyr Glu Thr Ser Phe Trp Leu

55



EP 1 043 395 A1

	355	360	365	
5	ctg tgg gcc aac tgg gcc gtc aac ccc gtc ctc tac cca ctg tgc cac			1152
	Leu Trp Ala Asn Ser Ala Val Asn Pro Val Leu Tyr Pro Leu Cys His			
	370	375	380	
10	tac agc ttc cgc aga gcc ttc acc aag ctc ctc tgc ccc cag aag ctc			1200
	Tyr Ser Phe Arg Arg Ala Phe Thr Lys Leu Leu Cys Pro Gln Lys Leu			
	385	390	395	400
15	aag gtc cag ccc cac gcc tcc ctg gag cag tgc tgg aag			1239
	Lys Val Gln Pro His Gly Ser Leu Glu Gln Cys Trp Lys			
	405	410		
20	<210> 3			
	<211> 21			
	<212> DNA			
	<213> Artificial Sequence			
25	<220>			
	<223> Description of Artificial Sequence: Artificially			
	synthesized primer sequence			
30	<400> 3			
	batngccaac ctbkcttct c			21
35	<210> 4			
	<211> 20			
	<212> DNA			
	<213> Artificial Sequence			
40	<220>			
	<223> Description of Artificial Sequence: Artificially			
	synthesized primer sequence			
45	<400> 4			
	ccataaaagn nggggttgac			20
50	<210> 5			
	<211> 2700			
	<212> DNA			
55				

EP 1 043 395 A1

<213> Rattus norvegicus

5 <220>  
 <221> CDS  
 <222> (351)..(1589)

10 <400> 5  
 aattcggcac gagcgggcag atcgcggggc gcaactcgggtt gcgcgctgag ctagggggtgc 60  
 accgacgcac cgcggggcggc tggagctcgg ctttgctctc gctgcagcag ccgcgcgcgc 120  
 15 cgccccactc cgctcagatt ccgacaccag cccctcttgg atcgccctcc tggactctag 180  
 cccgggctct tgctccgacc ccgcggacca tgctccgggc gccccccgga aaaccgggct 240  
 20 gggcgaagag ccggcaaaga ttaggtcac gagcgggggc cccaccggc caccagctc 300  
 tccgcccgtg cctgcccgg tgctcccgag ccgtgtgagc ctgctgggcc atg gag 356  
 25 Met Glu  
 1

cgc gcg ccg ccc gac ggg ctg atg aac gcg tcg ggc act ctg gcc gga 404  
 30 Arg Ala Pro Pro Asp Gly Leu Met Asn Ala Ser Gly Thr Leu Ala Gly  
 5 10 15

gag gcg gcg gct gca ggc ggg gcg cgc ggc ttc tcg gct gcc tgg acc 452  
 35 Glu Ala Ala Ala Ala Gly Gly Ala Arg Gly Phe Ser Ala Ala Trp Thr  
 20 25 30

gct gtc ctg gct gcg ctc atg gcg ctg ctc atc gtg gcc aca gta ctg 500  
 40 Ala Val Leu Ala Ala Leu Met Ala Leu Leu Ile Val Ala Thr Val Leu  
 35 40 45 50

ggc aac gcg ctg gtc atg ctc gcc ttc gtg gcg gat tcg agc ctc cgc 548  
 45 Gly Asn Ala Leu Val Met Leu Ala Phe Val Ala Asp Ser Ser Leu Arg  
 55 60 65

acc cag aac aac ttc ttt ctg ctc aac ctc gcc atc tcc gac ttc ctc 596  
 50 Thr Gln Asn Asn Phe Phe Leu Leu Asn Leu Ala Ile Ser Asp Phe Leu  
 70 75 80

55

EP 1 043 395 A1

5	gtg ggt gcc ttc tgc atc cca ttg tac gta ccc tat gtg ctg acc ggc Val Gly Ala Phe Cys Ile Pro Leu Tyr Val Pro Tyr Val Leu Thr Gly	644
	85 90 95	
10	cgt tgg acc ttc ggc cgg ggc ctc tgc aag ctg tgg ctg gtg gta gac Arg Trp Thr Phe Gly Arg Gly Leu Cys Lys Leu Trp Leu Val Val Asp	692
	100 105 110	
15	tac cta ctg tgt gcc tcc tcg gtc ttc aac atc gta ctc atc agc tat Tyr Leu Leu Cys Ala Ser Ser Val Phe Asn Ile Val Leu Ile Ser Tyr	740
	115 120 125 130	
20	gac cga ttc ctg tca gtc act cga gct gtc tcc tac agg gcc cag cag Asp Arg Phe Leu Ser Val Thr Arg Ala Val Ser Tyr Arg Ala Gln Gln	788
	135 140 145	
25	ggg gac acg aga cgg gcc gtt cgg aag atg gca ctg gtg tgg gtg ctg Gly Asp Thr Arg Arg Ala Val Arg Lys Met Ala Leu Val Trp Val Leu	836
	150 155 160	
30	gcc ttc ctg ctg tat ggg cct gcc atc ctg agt tgg gag tac ctg tct Ala Phe Leu Leu Tyr Gly Pro Ala Ile Leu Ser Trp Glu Tyr Leu Ser	884
	165 170 175	
35	ggt ggc agt tcc atc ccc gag ggc cac tgc tat gct gag ttc ttc tac Gly Gly Ser Ser Ile Pro Glu Gly His Cys Tyr Ala Glu Phe Phe Tyr	932
	180 185 190	
40	aac tgg tac ttt ctc atc acg gcc tcc acc ctc gag ttc ttc acg ccc Asn Trp Tyr Phe Leu Ile Thr Ala Ser Thr Leu Glu Phe Phe Thr Pro	980
	195 200 205 210	
45	ttc ctc agc gtt acc ttc ttc aac ctc agc atc tac ctg aac atc cag Phe Leu Ser Val Thr Phe Phe Asn Leu Ser Ile Tyr Leu Asn Ile Gln	1028
	215 220 225	
50	agg cgc acc cgc ctt cgg ctt gat ggg ggc cgt gag gct ggc cca gaa Arg Arg Thr Arg Leu Arg Leu Asp Gly Gly Arg Glu Ala Gly Pro Glu	1076
	230 235 240	
55	ccc cca cca gat gcc cag ccc tcg cca cct cca gct ccc ccc agc tgc	1124

EP 1 043 395 A1

	Pro Pro Pro Asp Ala Gln Pro Ser Pro Pro Pro Ala Pro Pro Ser Cys	
5	245 250 255	
	tgg ggc tgc tgg cca aaa ggg cat ggc gag gcc atg ccg ttg cac agc	1172
	Trp Gly Cys Trp Pro Lys Gly His Gly Glu Ala Met Pro Leu His Ser	
10	260 265 270	
	tct ggc agc tcc tca agg ggc act gag agg cca cgc tca ctc aaa agg	1220
	Ser Gly Ser Ser Ser Arg Gly Thr Glu Arg Pro Arg Ser Leu Lys Arg	
15	275 280 285 290	
	ggc tcc aag cca tca gca tct tca gca tcc ctg gag aag cgc atg aag	1268
	Gly Ser Lys Pro Ser Ala Ser Ser Ala Ser Leu Glu Lys Arg Met Lys	
20	295 300 305	
	atg gtg tcc cag agc atc acc cag cgc ttc cgg ctg tgc cgg gac aag	1316
	Met Val Ser Gln Ser Ile Thr Gln Arg Phe Arg Leu Ser Arg Asp Lys	
25	310 315 320	
	aag gtg gcc aag tgc ctg gcc atc atc gtg agc atc ttt ggg ctc tgc	1364
	Lys Val Ala Lys Ser Leu Ala Ile Ile Val Ser Ile Phe Gly Leu Cys	
30	325 330 335	
	tgg gcg ccg tac acg ctc cta atg atc atc cga gct gct tgc cat ggc	1412
	Trp Ala Pro Tyr Thr Leu Leu Met Ile Ile Arg Ala Ala Cys His Gly	
35	340 345 350	
	cgc tgc atc ccc gat tac tgg tac gag acg tcc ttc tgg ctt ctg tgg	1460
	Arg Cys Ile Pro Asp Tyr Trp TyrGlu Thr Ser Phe Trp Leu Leu Trp	
40	355 360 365 370	
	gcc aac tgc gcc gtc aac ccc gtc ctc tac cca ctg tgc cac tac agc	1508
	Ala Asn Ser Ala Val Asn Pro Val Leu Tyr Pro Leu Cys His Tyr Ser	
45	375 380 385	
	ttc cgc aga gcc ttc acc aag ctc ctc tgc ccc cag aag ctc aag gtc	1556
	Phe Arg Arg Ala Phe Thr Lys Leu Leu Cys Pro Gln Lys Leu Lys Val	
50	390 395 400	
	cag ccc cac ggc tcc ctg gag cag tgc tgg aag tgagcagctg cccaccctt	1609
	Gln Pro His Gly Ser Leu Glu Gln Cys Trp Lys	
55		

EP 1 043 395 A1

405

410

5 ctgaggccag gcccttgtae ttgtttgagt gggcagccgg agcgtgggag gggccctggt 1669

ccatgctccg ctccaaatgc catggcgccg tcttagatca tcaacccgcg agtggggtag 1729

10 catggcaggt gggccaagag ccctagttag tggagctaga gtgtgctggt tagctctgcc 1789

gccacattct ccttcaccac acagaagaga caatccagga gtcccaggca tgccttcac 1849

15 ctacacacac acacacacac acacacacac acacaccaca gtgcagtgcc agtgaatgcc 1909

ccttttgcat atttagtggc tgggtgcttc cctaatacaa accctgggtg gtgctcccg 1969

20 ctccggccct ggcaatgcgt gcgtgcgccc tgcattgtgt cacacccgcg acacacccgc 2029

ccgccacaca cttgcaacac ctcctctctc ccagaagagc tggggacgat gccctttgct 2089

25 gccactgtct cttgtttaat cccagagcct ggctccttat ccccaactct ccttcaact 2149

ctgccccaca aagtgtcgag cgctcggga aacttgaagc ttctctgctc ctccactct 2209

30 ggatgtttc aggaagatgg aggagaagaa aacacgtctg tgaacttgat gttccttgga 2269

tgtttaatca agagagacaa aattgccgag gagctcggg ctggattggc aggtgtggg 2329

35 tcccacgccc tctccctca gtgtgcagc ttcgggtga gccgcgccc agtctctgc 2389

ctgccccgc cccaggcttg ggacgatggc cctgccctgc ttgccccgc tgtacaatca 2449

40 gaatttggg gtgggtggtt atggggtaga gcggctcttc actgtgccct aaaggctctg 2509

aggctcacag gacagtcagc aggagagcag gcaggccgcg gacacctggg aggaatgctt 2569

45 tgctctgccc tgtgtactca cctcaggctt ctgcatgctc tgctgccctt gtgccctggt 2629

gtgctgcctc tgccaatgtg aaaacacaat aaagtgtatt tttttacgga aaaaaaana 2689

50 aaaaaaaaaa a 2700

55

10

15

29

20

25

30

35

40

45

50

1 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 10

&lt;220&gt;

<223> Description of Artificial Sequence: Artificially  
synthesized primer sequence

&lt;400&gt; 9

cttcgcgcgg gccttcacca a

21

&lt;210&gt; 10

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Artificially  
synthesized primer sequence

&lt;400&gt; 10

acagacacgg cggggctcac

20

&lt;210&gt; 11

&lt;211&gt; 1350

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; exon

&lt;222&gt; (280) .. (557)

&lt;400&gt; 11

gcactcggct gcgcgttgcn tccggctgca cggtcgcacc ggcagcggct caggctccgg 60

ctctctctcc gctgcagcag ccgcgctgcc ggccccactg ggctcggatc cggccccggc 120

cccctcggca ccgcctgctc tggccccggc cccggccccg cggaccatgc gctggggcgc 180

cccaggggaa cccgaccggg ccaaggggcc gcaaagacga ggctcccggg ccggggcccc 240

tcccgccgc ccagctctcg gccggcgccc tgccccgcgt cccggagccg cgtgagcctg 300

cggggccatg gagcgcgcgc cgccccacgg gccgctgaac gcttcggggg cgctggcggg 360

# EP 1 043 395 A1

cgaggcggcg gcggcggcg gggcgcgcg cttctcgca gcctggaccg cggtagctggc 420  
 5 cgcgtcatg gcgctgctca tcgtggccac ggtgctgggc aacgcgtgg tcattgctcgc 480  
 cttegtggcc gactcgagcc tccgcaccca gaacaacttc ttctgctca acctcgccat 540  
 10 ctccgacttc ctgctcggtta aatccccagc ccctggccgc tggggaccca gggcgccca 600  
 gcgtggccgg gccagcgggg actggaacac ggacctgggt ggctcccgca ggcacacgcc 660  
 15 ccaccagggg acccgccctg ggaagggggc gtccggagcc catggggtgg gggcacaggg 720  
 cgaagtccct tgccactcag gcctcgggac aggggctggg gagagatgtc cccgggaagg 780  
 20 gacacgggca ctggcgagg cgcaaggcgc aaaggcagcg ggtgcagctc tggtctctgc 840  
 gctgtagcca aacaaaggct gctgcggact taggacgcgc ggaggcgca gtggggcggt 900  
 25 ttagagaagg tctgggggag gggacatgga agggggattt ttagagctgt gttgggggaa 960  
 gggacggtgg ggaaggtggg ggttggggga gacgctcgga ggagcgtgct ctcacgtgtc 1020  
 30 caggctctgc tgccggctgg gggcgggggc acgcgagggg ggctggagcg ccagacacct 1080  
 gttggggctg tgaggtgctt ctcccagacg ctccaagccc gcttggcagt agtagtagcg 1140  
 35 gctggcggct ggcggctgca accaagtgcc ctttcagcca ggagaaaggc tttctccttg 1200  
 tctaagctga gaccgagggt tgtccagcgc cagggtaggg gctggagtcc agcgggggag 1260  
 40 gggagaagga aattgtcttc tttctcctt tgagggtggg gagggtgga cagaagtcca 1320  
 gggaatcccg actccaggct ctccggggtc 1350  
 45

<210> 12  
 <211> 448  
 50 <212> DNA  
 <213> Homo sapiens

55



&lt;220&gt;

&lt;221&gt; exon

&lt;222&gt; (259)..(425)

&lt;400&gt; 12

gagctcccca tgcctggatc atccctctcg cccccaggcc caggggacac agatagtgt 60

gggagctatg tgggggtgaa ggctggcggc agggcagagt ttgtggctga caccaggtgg 120

aggggtggtg agatgaggat ggctagtccc agaaaagcag ccaccatgtg accccaggtc 180

ccgcgggtgt ctgcgccttag gtccgtctgt cccctggccc ctggctgcat ggtcccaactg 240

tggccctact cccacagggc gccttctgca tcccactgta tgtacctac gtgctgacag 300

gccgctggac ctccggccgg ggctctgca agctgtggct ggtagtggac tacctgctgt 360

gcacctcttc tgccttcaac atcgtgtcga tcagctacga ccgcttctcg tcggtcaccc 420

gagcgggtgag tcctgggctg cggagctc 448

&lt;210&gt; 13

&lt;211&gt; 1893

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; exon

&lt;222&gt; (293)..(1209)

&lt;400&gt; 13

gagctcacag ctggtagggg gtggtaaaaca ggcagcctag cagagagtga gggttcaggt 60

tggccccagg gagcttctga ggctctcact gagtgtggca gggcaccagt ccgggacccc 120

agtggggagg gttagaggaa gggaggggaa agagggaggg agggaggaca ggaggggaaa 180

ggaggagcat tgctgctgag ggaagggccc acataggggc ccacaggcta cgggggacga 240

cccagcccaa tattctctcc gccccgccc tgaccagct gcccttctgc aggtctcata 300

EP 1 043 395 A1

ccggggccag caggggtgaca cggggcgggc agtgcggaag atgctgctgg tgtgggtgct 360  
5 ggcttctctg ctgtacggac cagccatcct gagctgggag tacctgtccg ggggcagctc 420  
catccccgag ggccactgct atgccgagtt cttctacaac tggtaactcc tcatacaggg 480  
10 ttccaccctg gagttcttta cggccttctc cagcgtcacc ttctttaacc tcagcatcta 540  
cctgaacatc cagagggcgca cccgcctccg gctggatggg gctcgagagg cagccggccc 600  
15 cgagccccct cccgaggccc agccctcacc acccccaccg cctggctgct ggggctgctg 660  
gcagaagggg cagggggagg ccatgccgct gcacaggatg ggggtgggtg agggggccgt 720  
20 aggcctctgag gccggggagg cgaccctcgg ggggtggcgt gggggcggtt ccgtggcttc 780  
accacctcc agctccggca gctcctcgag gggcactgag aggcgcgct cactcaagag 840  
25 gggctccaag ccacggcgt cctcgccctc actggagaag cgcatgaaga tgggtgccca 900  
gagcttcacc cagcgtttc ggctgtctcg ggacaggaaa gtggccaagt cgctggccgt 960  
30 catcgtgagc atctttgggc tctgctgggc ccatacacg ctgctgatga tcacccgggc 1020  
cgctgceat ggccactgcg tccctgacta ctggtacgaa acctcctct ggctcctgtg 1080  
35 ggccaactcg gctgtcaacc ctgtctctta cctctgtgc caccacagct tccgccgggc 1140  
cttcaccaag ctgctctgcc ccagaagct caaatccag cccacagct cctggagca 1200  
40 ctgcgggaag tgagtggccc accagagcct cctcagcca cgctctctc agcccaggtc 1260  
tctgggcat ctggccctgc tgccccctac ccggctcgtt ccccagggg tgagccccgc 1320  
45 cgtgtctgtg gccctctctt aatgccacgg cagccaccct gccatggagg cgcttctctg 1380  
ggctggccag agggccctc actggctgga ctggaggctg ggtggccggc cctgcccccc 1440  
50 acattctggc tccaccggga gggacagtct ggaggtccca gacatgctgc ccacccctg 1500  
55

ctggtgcecca cccttcgcag ttactggttg gtgtttctcc caaagcaagc acctgggtgt 1560  
 5 gctccaggct tcttgcccta gcagtttgcc tctgcacgtg cacacacctg cacacccctg 1620  
 cacacacctg cacaccgtcc ctctccccgg acaagcccag gacactgcct ttgctgcctt 1680  
 10 ctgtctcttg cataagcctc aggcctggcc ctttcacccc tcttcccacc aactctctct 1740  
 gcccccaaaa gtgtcaaggg gccctaggaa cctcgaagct gttctctgct tttccattct 1800  
 15 ggggtgttttc agaaagatga agaagaaaac atgtctgtga acttgatggt cctgggatgt 1860  
 ttaatcaaga gagacaaaat tgctgaggag etc 1893

20 <210> 14  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

25 <220>  
 <223> Description of Artificial Sequence: Artificially  
 synthesized primer sequence

30 <400> 14  
 tgaacgcttc gggggcgctg 20

35 <210> 15  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

40 <220>  
 <223> Description of Artificial Sequence: Artificially  
 synthesized primer sequence

45 <400> 15  
 gagatggcga gggtgagcag g 21

50 <210> 16  
 <211> 20  
 <212> DNA

55

5

15

25

30

35

40

45

50

55

DOCID: &lt;EP 1043395A1 | &gt;

&lt;400&gt; 19

tgcgcctctg gatgttcag

19

&lt;210&gt; 20

&lt;211&gt; 453

&lt;212&gt; PRT-

&lt;213&gt; Homo sapiens

&lt;400&gt; 20

Met Glu Arg Ala Pro Pro Asp Gly Pro Leu Asn Ala Ser Gly Ala Leu

1 5 10 15

Ala Gly Glu Ala Ala Ala Ala Gly Gly Ala Arg Gly Phe Ser Ala Ala

20 25 30

Trp Thr Ala Val Leu Ala Ala Leu Met Ala Leu Leu Ile Val Ala Thr

35 40 45

Val Leu Gly Asn Ala Leu Val Met Leu Ala Phe Val Ala Asp Ser Ser

50 55 60

Leu Arg Thr Gln Asn Asn Phe Phe Leu Leu Asn Leu Ala Ile Ser Asp

65 70 75 80

Phe Leu Val Gly Ala Phe Cys Ile Pro Leu Tyr Val Pro Tyr Val Leu

85 90 95

Thr Gly Arg Trp Thr Phe Gly Arg Gly Leu Cys Lys Leu Trp Leu Val

100 105 110

Val Asp Tyr Leu Leu Cys Thr Ser Ser Ala Phe Asn Ile Val Leu Ile

115 120 125

Ser Tyr Asp Arg Phe Leu Ser Val Thr Arg Ala Val Ser Tyr Arg Ala

130 135 140

Gln Gln Gly Asp Thr Arg Arg Ala Val Arg Lys Met Leu Leu Val Trp

145 150 155 160

Val Leu Ala Phe Leu Leu Tyr Gly Pro Ala Ile Leu Ser Trp Glu Tyr

# EP 1 043 395 A1

	165	170	175
5	Leu Ser Gly Gly Ser Ser Ile Pro Glu Gly His Cys Tyr Ala Glu Phe		
	180	185	190
10	Phe Tyr Asn Trp Tyr Phe Leu Ile Thr Ala Ser Thr Leu Glu Phe Phe		
	195	200	205
15	Thr Pro Phe Leu Ser Val Thr Phe Phe Asn Leu Ser Ile Tyr Leu Asn		
	210	215	220
20	Ile Gln Arg Arg Thr Arg Leu Arg Leu Asp Gly Ala Arg Glu Ala Ala		
	225	230	235
25	Gly Pro Glu Pro Pro Pro Glu Ala Gln Pro Ser Pro Pro Pro Pro		
	245	250	255
30	Gly Cys Trp Gly Cys Trp Gln Lys Gly His Gly Glu Ala Met Pro Leu		
	260	265	270
35	His Arg Tyr Gly Val Gly Glu Ala Ala Val Gly Ala Glu Ala Gly Glu		
	275	280	285
40	Ala Thr Leu Gly Gly Gly Gly Gly Gly Gly Ser Val Ala Ser Pro Thr		
	290	295	300
45	Ser Ser Ser Gly Ser Ser Ser Arg Gly Thr Glu Arg Pro Arg Ser Leu		
	305	310	315
50	Lys Arg Gly Ser Lys Pro Ser Ala Ser Ser Ala Ser Leu Glu Lys Arg		
	325	330	335
55	Met Lys Met Val Ser Gln Ser Phe Thr Gln Arg Phe Arg Leu Ser Arg		
	340	345	350
60	Asp Arg Lys Val Ala Lys Ser Leu Ala Val Ile Val Ser Ile Phe Gly		
	355	360	365
65	Leu Cys Trp Ala Pro Tyr Thr Leu Leu Met Ile Ile Arg Ala Ala Cys		
	370	375	380

EP 1043 395 A1

His Gly His Cys Val Pro Asp Tyr Trp Tyr Glu Thr Ser Phe Trp Leu  
 385 390 395 400

5  
 Leu Trp Ala Asn Ser Ala Val Asn Pro Val Leu Tyr Pro Leu Cys His  
 405 410 415

10  
 His Ser Phe Arg Arg Ala Phe Thr Lys Leu Leu Cys Pro Gln Lys Leu  
 420 425 430

15  
 Lys Ile Gln Pro His Ser Ser Leu Glu His Cys Trp Lys Lys Met Lys  
 435 440 445

Lys Lys Thr Cys Leu  
 450

20  
 <210> 21  
 <211> 2050  
 <212> DNA  
 25 <213> Homo sapiens

<220>  
 <221> CDS  
 30 <222> (271)..(1629)

<400> 21  
 agagatgtag ggcgccctt ttagctgcgc acagaacgaa agaactcggt ttttctttaa 60  
 35 gtagtggtgc ttgggtgacg cttagggcgc cctccgcagt gcgcgcagga aagcgactg 120

aggctgcgga ggcagagctg catgctgggt gcgggaagag gtgggctccg tcgcggagtc 180  
 40 gctgagtcgc tgccctttta gttagttctg cagtctagta tggccccat ttgcccttcc 240

actcccgag ccgcgtgagc ctgcggggcc atg gag cgc gcg ccg ccc gac ggg 294  
 45 Met Glu Arg Ala Pro Pro Asp Gly  
 1 5

ccg ctg aac gct tcg ggg gcg ctg gcg ggc gag gcg gcg gcg gcg ggc 342  
 50 Pro Leu Asn Ala Ser Gly Ala Leu Ala Gly Glu Ala Ala Ala Ala Gly  
 10 15 20

55

# EP 1 043 395 A1

	ggg gcg cgc ggc ttc tgc gca gcc tgg acc gcg gtg ctg gcc gcg ctc	390
	Gly Ala Arg Gly Phe Ser Ala Ala Trp Thr Ala Val Leu Ala Ala Leu	
5	25 30 35 40	
	atg gcg ctg ctc atc gtg gcc acg gtg ctg ggc aac gcg ctg gtc atg	438
	Met Ala Leu Leu Ile Val Ala Thr Val Leu Gly Asn Ala Leu Val Met	
10	45 50 55	
	ctc gcc ttc gtg gcc gac tgc agc ctc cgc acc-cag aac aac ttc ttc	486
	Leu Ala Phe Val Ala Asp Ser Ser Leu Arg Thr Gln Asn Asn Phe Phe	
15	60 65 70	
	ctg ctc aac ctc gcc atc tcc gac ttc ctc gtc ggc gcc ttc tgc atc	534
	Leu Leu Asn Leu Ala Ile Ser Asp Phe Leu Val Gly Ala Phe Cys Ile	
20	75 80 85	
	cca ctg tat gta ccc tac gtg ctg aca gcc cgc tgg acc ttc ggc cgg	582
	Pro Leu Tyr Val Pro Tyr Val Leu Thr Gly Arg Trp Thr Phe Gly Arg	
25	90 95 100	
	ggc ctc tgc aag ctg tgg ctg gta gtg gac tac ctg ctg tgc acc tcc	630
	Gly Leu Cys Lys Leu Trp Leu Val Val Asp Tyr Leu Leu Cys Thr Ser	
30	105 110 115 120	
	tct gcc ttc aac atc gtg ctc atc agc tac gac cgc ttc ctg tgc gtc	678
	Ser Ala Phe Asn Ile Val Leu Ile Ser Tyr Asp Arg Phe Leu Ser Val	
35	125 130 135	
	acc cga gcg gtc tca tac cgg gcc cag cag ggt gac acg cgg cgg gca	726
	Thr Arg Ala Val Ser Tyr Arg Ala Gln Gln Gly Asp Thr Arg Arg Ala	
40	140 145 150	
	gtg cgg aag atg ctg ctg gtg tgg gtg ctg gcc ttc ctg ctg tac gga	774
	Val Arg Lys Met Leu Leu Val Trp Val Leu Ala Phe Leu Leu Tyr Gly	
45	155 160 165	
	cca gcc atc ctg agc tgg gag tac ctg tcc ggg gcc agc tcc atc ccc	822
	Pro Ala Ile Leu Ser Trp Glu Tyr Leu Ser Gly Gly Ser Ser Ile Pro	
50	170 175 180	
	gag gcc cac tgc tat gcc gag ttc ttc tac aac tgg tac ttc ctc atc	870
55		



EP 1 043 395 A1

Glu Gly His Cys Tyr Ala Glu Phe Phe Tyr Asn Trp Tyr Phe Leu Ile  
 185 190 195 200  
 5  
 arg gct tcc acc ctg gag ttc ttt acg ccc ttc ctc agc gtc acc ttc 918  
 Thr Ala Ser Thr Leu Glu Phe Phe Thr Pro Phe Leu Ser Val Thr Phe  
 10 205 210 215  
 ttt aac ctc agc atc tac ctg aac atc cag agg cgc acc cgc ctc cgg 966  
 Phe Asn Leu Ser Ile Tyr Leu Asn Ile Gln Arg Arg Thr Arg Leu Arg  
 15 220 225 230  
 ctg gat ggg gct cga gag gca gcc ggc ccc gag ccc cct ccc gag gcc 1014  
 Leu Asp Gly Ala Arg Glu Ala Ala Gly Pro Glu Pro Pro Pro Glu Ala  
 20 235 240 245  
 cag ccc tca cca ccc cca ccg cct ggc tgc tgg ggc tgc tgg cag aag 1062  
 Gln Pro Ser Pro Pro Pro Pro Gly Cys Trp Gly Cys Trp Gln Lys  
 25 250 255 260  
 ggg cac ggg gag gcc atg ccg ctg cac agg tat ggg gtg ggt gag gcg 1110  
 Gly His Gly Glu Ala Met Pro Leu His Arg Tyr Gly Val Gly Glu Ala  
 30 265 270 275 280  
 gcc gta ggc gct gag gcc ggg gag gcg acc ctc ggg ggt ggc ggt ggg 1158  
 Ala Val Gly Ala Glu Ala Gly Glu Ala Thr Leu Gly Gly Gly Gly Gly  
 35 285 290 295  
 ggc ggc tcc gtg gct tca ccc acc tcc agc tcc ggc agc tcc tcg agg 1206  
 Gly Gly Ser Val Ala Ser Pro Thr Ser Ser Ser Gly Ser Ser Ser Arg  
 40 300 305 310  
 ggc act gag agg ccg cgc tca ctc aag agg ggc tcc aag ccg tcg gcg 1254  
 Gly Thr Glu Arg Pro Arg Ser Leu Lys Arg Gly Ser Lys Pro Ser Ala  
 45 315 320 325  
 tcc tcg gcc tcg ctg gag aag cgc atg aag atg gtg tcc cag agc ttc 1302  
 Ser Ser Ala Ser Leu Glu Lys Arg Met Lys Met Val Ser Gln Ser Phe  
 50 330 335 340  
 acc cag cgc ttt cgg ctg tct cgg gac agg aaa gtg gcc aag tcg ctg 1350  
 Thr Gln Arg Phe Arg Leu Ser Arg Asp Arg Lys Val Ala Lys Ser Leu  
 55

EP 1043 395 A1

	345	350	355	360	
5	gcc gtc atc gtg agc atc ttt ggg ctc tgc tgg gcc cca tac acg ctg				1398
	Ala Val Ile Val Ser Ile Phe Gly Leu Cys Trp Ala Pro Tyr Thr Leu				
	365		370	375	
10	ctg atg atc atc cgg gcc gcc tgc cat ggc cac tgc gtc cct gac tac				1446
	Leu Met Ile Ile Arg Ala Ala Cys His Gly His Cys Val Pro Asp Tyr				
	380		385	390	
15	tgg tac gaa acc tcc ttc tgg ctc ctg tgg gcc aac tgc gct gtc aac				1494
	Trp Tyr Glu Thr Ser Phe Trp Leu Leu Trp Ala Asn Ser Ala Val Asn				
	395		400	405	
20	cct gtc ctc tac cct ctg tgc cac cac agc ttc cgc cgg gcc ttc acc				1542
	Pro Val Leu Tyr Pro Leu Cys His His Ser Phe Arg Arg Ala Phe Thr				
	410		415	420	
25	aag ctg ctc tgc ccc cag aag ctc aaa atc cag ccc cac agc tcc ctg				1590
	Lys Leu Leu Cys Pro Gln Lys Leu Lys Ile Gln Pro His Ser Ser Leu				
	425		430	435	440
30	gag cac tgc tgg aaa aag atg aag aag aaa aca tgt ctg tgaacttgat				1639
	Glu His Cys Trp Lys Lys Met Lys Lys Lys Thr Cys Leu				
	445		450		
35	gttcctggga tgcttaataca agagagacaa aattgctgag gagctcaggg ctggattggc				1699
	agggtgtgggc tcccacgccc tcttccctcc gctaaggctt ccggctgagc tgtgccagct				1759
40	gcttctgccc accccgcttc tgggtcaca ccagccctgg tggccaagcc tgccccggcc				1819
	actctgtttg ctcacccagg acctcgggg gtgtgtggga ggagggggcc cggctgggcc				1879
45	cgagggtccc aaggcgtgca ggggcggtcc agaggaggtg cccgggcagg ggccgcttcg				1939
	ccatgtgctg tgcacccgtg ccacgcctc tgcctgctcc tctgctctg cccgctgcgc				1999
50	tgccctgcaa accgtgaggt cacaataaag tgtatttttt tattggtgct g				2050
	<210> 22				

55

5 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 10 <220>  
 <223> Description of Artificial Sequence: Artificially  
 synthesized primer sequence  
  
 15 <400> 22  
 tgcaccccat tgtacgtacc 20  
  
 20 <210> 23  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 25 <220>  
 <223> Description of Artificial Sequence: Artificially  
 synthesized primer sequence  
  
 30 <400> 23  
 atcattagga gcgtgtangg 20  
  
 35 <210> 24  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence  
  
 40 <220>  
 <223> Description of Artificial Sequence: Artificially  
 synthesized primer sequence  
  
 45 <400> 24  
 tgcctcggga caccatcttc 20  
  
 50  
 1  
  
 55

Claims

1. A guanosine triphosphate binding protein-coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 1, or said amino acid sequence in which one or more amino acids are replaced, deleted, or added.
2. A guanosine triphosphate binding protein-coupled receptor protein comprising the amino acid sequence of SEQ ID NO: 20, or said amino acid sequence in which one or more amino acids are replaced, deleted, or added.
3. A guanosine triphosphate binding protein-coupled receptor protein encoded by a DNA which hybridizes with a DNA comprising the nucleotide sequence of SEQ ID NO: 2.
4. A guanosine triphosphate binding protein-coupled receptor protein encoded by a DNA which hybridizes with a DNA comprising the nucleotide sequence of SEQ ID NO: 21.
5. A partial peptide of the receptor protein as described in any one of claims 1 to 4.
6. A DNA encoding the receptor protein as described in any one of claims 1 to 4 or the partial peptide as described in claim 5.
7. The DNA described in claim 6, wherein said DNA comprising the nucleotide sequence of SEQ ID NO: 2 or NO: 21.
8. A vector containing the DNA as described in any one of claims 5 to 7.
9. A transformant carrying the vector as described in claim 8.
10. A method of producing the receptor protein as described in any one of claims 1 to 4 or the partial peptide as described in claim 5, the method comprising culturing the transformant as described in claim 9.
11. A method of screening for a ligand of the receptor protein as described in any one of claims 1 to 4, the method comprising exposing a test compound to the receptor protein as described in any one of claims 1 to 4 or the partial peptide as described in claim 5, and selecting a compound that binds to said protein or partial peptide.
12. A method of screening for a compound that inhibits the binding between the receptor protein as described in any one of claims 1 to 4 and its ligand, the method comprising,
  - (a) exposing a ligand to the receptor protein as described in any one of claims 1 to 4 or the partial peptide as described in claim 5 in the presence of a test compound, and detecting the binding activity between said protein or partial peptide and the ligand, and,
  - (b) comparing the binding activity detected in (a) with that in the absence of the test compound, and selecting a compound that reduces the binding activity between said protein or partial peptide and the ligand.
13. A kit for screening a compound that inhibits the binding between the receptor protein as described in any one of claims 1 to 4 and its ligand, the kit comprising the receptor protein as described in any of claims 1 to 4 or the partial peptide as described in claim 5.
14. An antibody which binds to the receptor protein as described in any one of claims 1 to 4.

Figure 1




Figure 2






Figure 3



**Figure 4**





## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP98/05967

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> Int.Cl <sup>1</sup> C12N15/12, C12N15/63, C12Q1/02, C12P21/08, C07K14/705, C07K16/28, C12N1/21, C12N1/19, C12N5/12 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) Int.Cl <sup>1</sup> C12N15/12, C12N15/63, C12Q1/02, C12P21/08, C07K14/705, C07K16/28, C12N1/21, C12N1/19, C12N5/12 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WPI (DIALOG), BIOSIS (DIALOG), EMBL/DDBJ/GenBank/Geneseq		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Bonner, T.I. et al., "Identification of a Family of Muscarinic Acetylcholine Receptor Genes" Science (1987) Vol. 237 p.527-532	1, 3, 5-14
Y	Dewan, Z. et al., "Molecular characterization of a rat arufa2B-adrenergic receptor" Proc. Natl. Acad. Sci. USA (1990) Vol. 87, No. 8 p.3102-3106	1, 3, 5-14
Y	JP, 7-67654, A (Mitsubishi Kagaku Bio-Chemical Laboratories, Inc.), 14 March, 1995 (14. 03. 95) (Family: none)	2, 4, 5-14
Y	Ernest, G.P. et al., "Distinct primary structures, ligand-binding properties and tissue-specific expression of four human muscarinic acetylcholine receptors" The ENBO J. (1987) Vol. 6, No. 13 p.3923-3929	2, 4-14
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 12 March, 1999 (12. 03. 99)		Date of mailing of the international search report 30 March, 1999 (30. 03. 99)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/JP98/05967

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Thomas, B. et al., "A Novel Subtype of Muscarinic Receptor Identified by Homology Screening" Biochim. Biophys. Res. Commun. (1987) Vol. 149, No. 1 p.125-132	1-14
A	Maria, F.B. et al., "Normalization and Subtraction: Two Approaches to Facilitate Gene Discovery" Genome Res. (1996) Vol. 6, No. 9 p.791-806	1-14

Form PCT/ISA/210 (continuation of second sheet) (July 1992)